



Thyroid Disease

Purpose

To identify and promote the most important contributors to improved clinical outcomes for thyroid disease.

Key Recommendations

- Consider patients for evaluation for hypothyroidism and hyperthyroidism every 5 years as part of a physical or health maintenance visit.
- Order TSH for patients with signs, symptoms or other indications of hypothyroidism. Order Free T4 for those with pituitary disease, brain injury or in the first trimester of pregnancy.
- Order TSH for suspected hyperthyroidism and confirm elevated FT4 and/or FT3.
- Become familiar with the key points about thyroid nodules.

High Risk Populations/Disparities

- Women of Jewish descent are nearly two times more likely than other women to have thyroid disease caused by an overactive immune system.¹
- Among patients with thyroid cancer, blacks have lower survival rates at five years than do whites. A study of a large cancer database found that blacks are 2.3 times more likely to get the most aggressive form of thyroid cancer (anaplastic). Also, thyroid tumors more often are larger and found at a later stage in blacks.¹
- In the National Health and Nutrition Examination Survey (NHANES), among individuals with no thyroid disease, hypothyroidism and hyperthyroidism was significantly higher in females than males.²
- BMI is positively associated with thyroid cancer risk in both men and women.³

Thyroid Disease

Identification of Thyroid Disease

Patients should be considered for evaluation for hyperthyroidism and hypothyroidism every 5 years as part of a physical or health maintenance visit.

Case Finding:

- Order TSH for patients with signs, symptoms, or other indications of hypo- or hyperthyroidism.
- Other indications, patients:
 - with autoimmune diseases (e.g. Type 1 diabetes mellitus, B12 deficiency, pernicious anemia, Addison's disease and collagen-vascular diseases)
 - with previous thyroid injury (exposure to radiation, excess iodine)
 - with previous thyroid surgery or thyroid function abnormality
 - when patient is undergoing treatment with interferon for hepatitis C perform baseline testing before treatment and every 3 months during treatment
 - when patient is undergoing treatment with amiodarone or lithium perform baseline testing before treatment and every 3 months during treatment

Testing:

- TSH is the best choice except in rare cases
- TSH + FT4 for patients with secondary diagnosis of thyroid disease and other CNS diseases, including brain injury

Refer to endocrinologist if results of tests confusing

Identification of Thyroid Disease for Pregnant Women and Those Planning Pregnancy:

- At initial interview, ask questions to identify potential thyroid problems and test as appropriate. Factors to be considered in case finding:
 - Those with autoimmune diseases often associated with thyroid disease, such as type 1 diabetes and pernicious anemia
 - Patients with a prior history of thyroid disease or thyroid surgery, an abnormal thyroid exam, or taking drugs known to affect the thyroid
 - Patients with a family history of thyroid illness or history of miscarriage
- When testing needed, first trimester, perform TSH and FT4
- This is necessary since in the first trimester HCG mimics TSH and leads to suppressed TSH even when patient euthyroid
- Second/third trimester perform TSH except when thyroid disease is secondary diagnosis (cause is hypothalamic or pituitary)

Treatment of Pregnant Women with Thyroid Disease

- Pregnant women who have hypothyroidism and those on thyroxine, test on regular basis (see above) during pregnancy, treat appropriately
- Thyroid nodules can and should be evaluated, follow up as necessary with obstetrician/gynecologist, surgeon & endocrinologist working together
- Imaging (ultrasound) may be ordered, as indicated, **radioisotope studies should not be done**

Hypothyroidism

Hypothyroidism is a syndrome that results from inadequate levels of thyroid hormone (T-4 and/or T-3). It is manifested by a set of symptoms, physical findings and laboratory tests.

See pages 3-5 for algorithms on diagnosis and treatment of hypothyroidism and hyperthyroidism and thyroid nodule

Key Points about Hypothyroidism

- Signs and symptoms of hypothyroidism: decrease in metabolic rate, tiredness, lethargy, sensitivity to cold, menstrual disturbances, goiter, hyperlipidemia
- **For most patients, TSH is the test of choice** for screening and monitoring of hypothyroidism
- Numerous drugs/diseases can alter T4 & T3 levels
- TSH is *not* the test of choice for those with pituitary disease, brain injury or in first trimester of pregnancy
- What is normal TSH (within the reference range) depends largely on individual

Key Points about Hyperthyroidism

- TSH the best diagnostic test for hyperthyroidism
- Asymptomatic patients over age 50 with suppressed TSH (less than 0.1) at risk for atrial fibrillation
- Confirmation of elevated FT4 and/or FT3 required after screening with a TSH
- Workup should include iodine uptake and scan to define etiology of hyperthyroidism
- Review treatment options with patient and family

Key Points about Thyroid Nodule (TN) Evaluation

- Rule of FIVE: 5% of your overall patient population will have palpable TNs, 5% of all palpable TNs are malignant
- Rule of ACTIVITY: Most nodules that are hyperactive are benign (extremely rare exceptions)

Refer to Endocrinologist

- RISK FACTORS for malignancy:
 - A PALPABLE nodule
 - Age < 20 or >60
 - M>F
 - History of radiation exposure
 - Familial history of thyroid cancer
 - Progressive growth on oral T-4
 - Ultrasound characteristics of TN causing concern:
 - Microcalcifications
 - Hypoechogenicity
 - Intranodule vascularity
 - Irregular nodule margins
 - Nodule > 10mm
 - There is no strong evidence that treatment of euthyroid patients with thyroid hormone will shrink benign thyroid nodules.
 - Evaluate further if patient has normal or elevated TSH
 - Evaluate all palpable nodules
- See endocrinologist for any nodule with above risk factors or concerning

Thyroid Disease




Figure 1. Treatment Algorithm for Subclinical Hypothyroidism.

This algorithm does not apply to pregnant women or to young women who may potentially seek pregnancy and is based on current U.S. and European guidelines.^{26,27} However, the U.S. guidelines do not make an explicit distinction according to age, and neither guideline specifies differential management according to the degree of thyrotropin elevation below 10 mIU per liter.

There are special considerations from pregnancy, both testing and replacement.

From New England Journal of Medicine, Robin P. Peeters, M.D., Ph.D, Subclinical Hypothyroidism, Volume No. 376, Page No. 2561. Copyright © 2017 Massachusetts Medical Society. Reprinted with permission from Massachusetts Medical Society.

Guidelines are intended to be flexible. They serve as reference points or recommendations, not rigid criteria. Guidelines should be followed in most cases, but there is an understanding that, depending on the patient, the setting, the circumstances, or other factors, care can and should be tailored to fit individual needs.

Thyroid Disease

Thyroid Nodule Evaluation

Thyroid nodule <1cm, repeat in 1 yr. Consider fine needle aspiration (FNA) if nodule demonstrates listed characteristics of concern.

Thyroid Nodule (TN) Evaluation


1. Rule of FIVE: 5% of your overall patient population will have palpable TNs. 5% of all palpable TNs are malignant.
2. Rule of ACTIVITY: All nodules (extremely rare exceptions) that are hyperactive are benign.
3. RISK FACTORS for malignancy:
 - A palpable nodule
 - Age <20 or >60
 - M > F
 - History of radiation exposure
 - Familial history of thyroid cancer
 - Progressive growth on oral T-4

Ultrasound characteristics that cause concern with thyroid nodules:

1. Microcalcifications
2. Hypochoogenicity
3. Intranodule vascularity
4. Irregular nodule margins
5. Nodule >10mm

NOTES:

- Evaluate further if patient has normal or elevated TSH.
- All palpable nodules should be evaluated.
- Nodule <1cm with abnormal ultrasound features and multinodular goiters, consider endocrine evaluation.



Thyroid Disease

Summary of Recommendations for Thyroid Testing from National Organizations

Organization	Core Recommendation	Recommendation – Pregnant Women
American Academy of Family Physicians	Evidence is insufficient to recommend for or against routine screening for thyroid disease in adults. <i>(Clinical Considerations: http://www.aafp.org/patient-care/clinical-recommendations/all/thyroid.html)</i>	
American Association of Clinical Endocrinologists (AACE) and American Thyroid Association (ATA) Taskforce on Hypothyroidism in Adults	AACE recommends routine TSH measurement in older patients – age not specified – especially women. ATA recommends screening at age 35 yrs and every 5 yrs thereafter. <i>(Jeffrey R. Garber, Rhoda H. Cobin, Hossein Gharib, James V. Hennessey, Irwin Klein, Jeffrey I. Mechanick, Rachel Pessah-Pollack, Peter A. Singer, and Kenneth A. Woeber for the American Association of Clinical Endocrinologists and American Thyroid Association Taskforce on Hypothyroidism in Adults. Thyroid. December 2012, 22(12): 1200-1235. doi:10.1089/thy.2012.0205)</i>	It remains unclear if screening for hypothyroidism in pregnancy is beneficial. A consensus statement in 2004 and clinical practice guidelines in 2007 and 2011 found insufficient data to support a 1999 and 2005 recommendation for universal screening for thyroid dysfunction during pregnancy, but rather recommended aggressive case finding. <i>(Jeffrey R. Garber, Rhoda H. Cobin, Hossein Gharib, James V. Hennessey, Irwin Klein, Jeffrey I. Mechanick, Rachel Pessah-Pollack, Peter A. Singer, and Kenneth A. Woeber for the American Association of Clinical Endocrinologists and American Thyroid Association Taskforce on Hypothyroidism in Adults. Thyroid. December 2012, 22(12): 1200-1235. doi:10.1089/thy.2012.0205)</i>
American College of Obstetrics and Gynecologists		Thyroid testing is not a routine part of prenatal care and should be limited to women with symptoms of thyroid disease and those with a history of thyroid disease or other medical conditions associated with it, such as diabetes. <i>(Committee Opinion #381, Subclinical Hypothyroidism in Pregnancy, Obstetrics & Gynecology. October 2007. Practice Bulletin Number 148: Thyroid disease in pregnancy, April 2015. Obstet Gynecol. 2015;125: 996–1005 - recommendation to not perform routine screening for thyroid disease in pregnancy has not changed.)</i>
The Endocrine Society		The committee could not reach agreement regarding screening recommendations for all newly pregnant women. 1) Some members recommended screening of all pregnant women for serum TSH abnormalities by the ninth week or at the time of their first visit. 2) Some members recommended neither for nor against universal screening of all pregnant women for TSH abnormalities at the time of their first visit. These members strongly support aggressive case finding to identify and test high-risk women (Table 1) for elevated TSH concentrations by the ninth week or at the time of their first visit before and during pregnancy. <i>(Abalovich M, Alexander EK, Amino N, Barbour L, Cobin RH, De Groot L, et.al. Management of Thyroid Dysfunction during Pregnancy and Postpartum: An Endocrine Society Clinical Practice Guideline. J Clin Endocrinol Metab, August 2012, 97(8):2543–2565.)</i>
United States Preventive Services Task Force	Current evidence is insufficient to assess the balance of benefits and harms of screening for thyroid dysfunction in nonpregnant, asymptomatic adults. <i>(Released March 2015.)</i>	

Guidelines are intended to be flexible. They serve as reference points or recommendations, not rigid criteria. Guidelines should be followed in most cases, but there is an understanding that, depending on the patient, the setting, the circumstances, or other factors, care can and should be tailored to fit individual needs.



Thyroid Disease

Resources for Patients

American Thyroid Association

- Patients and Public – Thyroid Information - Provides up-to-date online information for thyroid patients and their families and free patient brochures about thyroid disease.
- Complementary and Alternative Medicine (CAM) - Provides information about thyroid disease and CAM, reviews types of CAM and stresses importance of patient's talking to their physician before taking CAM.

Hormone Health Network

Provides online information about hypothyroidism, hyperthyroidism, thyroid nodules and thyroid disorder symptoms.

National Institute of Diabetes, Digestive and Kidney Diseases

Provides online information about thyroid and testing physicians may order.

Thyroid Disease

References

1. Health Disparities in Hormone Disorders. Hormone Health Network. October 2013. Available at: www.hormone.org/hormones-and-health/scientific-statements/health-disparities
 2. Golden SH, Brown A, Cauley JA, Chin MA, Gary-Webb TL, Kim C, et.al. Health Disparities in Endocrine Disorders: Biological, Clinical, and Nonclinical Factors – An Endocrine Society Scientific Statement. The Journal of Clinical Endocrinology and Metabolism. 2012; 97(9): E1579-E1639. doi: 10.1210/jc.2012-2043. Available at: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3431576/>
 3. Kitahara CM, Platz EA, Freeman LB, Hsing AW, Linet MS, Park Y, et. al. Obesity and thyroid cancer risk among U.S. men and women: a pooled analysis of 5 prospective studies. Cancer Epidemiology, Biomarkers and Prevention. A publication of the American Association for Cancer Research; cosponsored by the American Society of Preventive Oncology. 2011; 20(3): 464-472. doi: 10.1158/1055-9965.EPI-10-1220. Available at: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3079276/>
- Abalovich M, Amino N, Barbour LA, Cobin RH, De Groot LJ, Glinoer D, Mandel SJ, Stagnaro-Green A. Management of Thyroid Dysfunction During Pregnancy and Postpartum: an Endocrine Society Clinical Practice Guideline. J Clin Endocrinol Metab 2007 Aug;92(8 Suppl):S1–S47. Available at: http://press.endocrine.org/doi/10.1210/jc.2007-0141?url_ver=Z39.88-2003&rfr_id=ori:rid:crossref.org&rfr_dat=cr_pub%3dpubmed
- American College of Obstetricians and Gynecologists. ACOG Committee Opinion No. 381. Subclinical Hypothyroidism in Pregnancy. Obstet Gynecol 2007;110:959–60. Available at: <http://www.acog.org/About-ACOG/News-Room/News-Releases/2007/Routine-Thyroid-Screening-Not-Recommended-for-Pregnant-Women>
- American College of Obstetricians and Gynecologists. ACOG Practice Bulletin Number 148: Thyroid disease in pregnancy. April 2015. Obstet Gynecol. 2015;125: 996–1005. Available at: <http://www.acog.org/Search?Keyword=thyroid+disease+in+pregnancy>
- David S. Cooper, Gerard M. Doherty, Bryan R. Haugen, Richard T. Kloos, Stephanie L. Lee, Susan J. Mandel, Ernest L. Mazzaferri, Bryan McIver, Furio Pacini, Martin Schlumberger, Steven I. Sherman, David L. Steward, and R. Michael Tuttle. Revised American Thyroid Association Management Guidelines for Patients with Thyroid Nodules and Differentiated Thyroid Cancer: The American Thyroid Association (ATA) Guidelines Taskforce on Thyroid Nodules and Differentiated Thyroid Cancer. Thyroid. November 2009, 19(11): 1167-1214. Available at: <http://deepblue.lib.umich.edu/bitstream/handle/2027.42/78131/thy.2009.0110.pdf?sequence=1&isAllowed=y>
- Jeffrey R. Garber, Rhoda H. Cobin, Hossein Gharib, James V. Hennessey, Irwin Klein, Jeffrey I. Mechanick, Rachel Pessah-Pollack, Peter A. Singer, and Kenneth A. Woeber for the American Association of Clinical Endocrinologists and American Thyroid Association Taskforce on Hypothyroidism in Adults. Thyroid. December 2012, 22(12): 1200-1235. Available at: <http://online.liebertpub.com/doi/abs/10.1089/thy.2012.0205>
- The American Thyroid Association Taskforce on Thyroid Disease During Pregnancy and Postpartum, Alex Stagnaro-Green, Marcos Abalovich, Erik Alexander, Fereidoun Azizi, Jorge Mestman, Roberto Negro, Angelita Nixon, Elizabeth N. Pearce, Offie P. Soldin, Scott Sullivan, and Wilmar Wiersinga. Thyroid. October 2011, 21(10): 1081-1125. doi:10.1089/thy.2011.0087. Available at: <http://online.liebertpub.com/doi/full/10.1089/thy.2011.0087>
- The American Thyroid Association and American Association of Clinical Endocrinologists Taskforce on Hyperthyroidism and Other Causes of Thyrotoxicosis, Rebecca S. Bahn, Henry B. Burch, David S. Cooper, Jeffrey R. Garber, M. Carol Greenlee, Irwin Klein, Peter Laurberg, I. Ross McDougall, Victor M. Montori, Scott A. Rivkees, Douglas S. Ross, Julie Ann Sosa, and Marius N. Stan. Thyroid. June 2011, 21(6): 593-646. Available at: <http://online.liebertpub.com/doi/abs/10.1089/thy.2010.0417>
- United States Preventive Services Task Force. Thyroid Dysfunction Screening. March 2015. Available at: www.uspreventiveservicestaskforce.org/Page/Document/UpdateSummaryFinal/thyroid-dysfunction-screening