



Vitamin D Screening Recommendations

Purpose

To provide guidance regarding population-based screening for vitamin D deficiency.

Key Messages

- Do not screen for vitamin D deficiency in healthy adults or children. Consistent with the recommendation of the American Society of Clinical Pathology as part of the *Choosing Wisely Program*, the New York State Department of Health has informed NY State Health Plans that population based screening (routine screening of otherwise healthy adults and children) for 25-OH-Vitamin D Deficiency should not be performed and that such screening should not be reimbursed under Medicaid.
- Nearly all Americans and Canadians obtain sufficient vitamin D from their diet.
- On a case-by-case basis, consider selective testing for vitamin D deficiency in high-risk individuals such as patients with malabsorption syndromes or osteoporosis and elderly (i.e. institutionalized) and infants with low sunlight exposure.

Overview

Numerous observational studies have shown a correlation between vitamin D and bone health, cardiovascular disease, cancer, diabetes, immune function, renal disease and stroke. Randomized controlled studies have shown that vitamin D is important for bone health, but they have not established causal relationship between vitamin D and other health outcomes, including mental health.

The Institute of Medicine (IOM), US Preventive Services Task Force (USPSTF) and the American Society of Clinical Pathology as part of the Choosing Wisely® campaign do not recommend screening for vitamin D deficiency in healthy individuals.

Establishing the cut point for the diagnosis of vitamin D deficiency is controversial because of the lack of standardization among assays and because some groups of patients, e.g. obese and African Americans, tend to have low levels of 25-Hydroxyvitamin D but have lower risk of fracture than the general population. This may be due to increased bioavailable vitamin D for which there is not currently commercially available testing. Lack of a cut point that is clear and consistent across labs has resulted in over diagnosis and encouraged excessive vitamin D testing. The IOM recommended a cut point of 20 ng/mL of serum 25-Hydroxyvitamin D for the diagnosis of vitamin D deficiency, indicating that levels above greater than or equal to 20 ng/mL are sufficient for 97.5% of the population.

The IOM, USPSTF and the Endocrine Society, state that the levels of vitamin D intake listed below sufficient for healthy individuals:

Age	Recommended Vitamin D Intake (IU/day)
0-12 months	400 IU
1-70 years	600 IU
>70 years	800 IU



Vitamin D Screening Recommendations

References

- Chung M, Balk EM et al. Vitamin D and Calcium: A Systematic Review of Health Outcomes. Evidence Report No. 183 (Prepared by the Tufts Evidence-based Practice Center under the Agency for Healthcare Research and Quality) August 2009.
- Dawson-Hughes B, Mithal A et al. International Osteoporosis Foundation Position Statement: Vitamin D Recommendations for Older Adults. *Osteoporosis International* July 2010 21:7 1151-1154.
- Judd S MPH PhD Tangpricha V MD PhD Vitamin D Deficiency and Risk for Cardiovascular Disease Symposium Article. *The American Journal of the Medical Services* July 2009 338:1 40-44.
- Kulie, T MD, Groff, A DO et al. Vitamin D: An Evidence-Based Review. *Journal of the American Board of Family Medicine* Nov. - Dec. 2009 22:6 698-706.
- LeFevre, Michael L. MD MSPH Screening for Vitamin D Deficiency in Adults: U.S. Preventive Services Task Force Recommendation Statement. *Annals of Internal Medicine* January 20, 2015 Vol. 162 No. 2 133-141
- Llewellyn, D PhD, Lang, I. PhD et al. Vitamin D and Risk of Cognitive Decline in Elderly Persons. *Archives of Internal Medicine* July 12, 2010 170:1.3.
- National Institute for Health and Clinical Excellence Antenatal Care Routine Care for the Healthy Pregnant Woman March 2008.
- National Cancer Institute Vitamin D and Cancer Prevention: Strengths and Limits of the Evidence. U.S. Department of Health and Human Services National Institute of Health 2009.
- Office of Dietary Supplements National Institutes of Health Dietary Supplement Fact Sheet: Vitamin D Health Professional Fact Sheet updated November 10, 2014 <http://ods.od.nih.gov/factsheets/VitaminD.asp>.
- Ontario Health Technology Advisory Committee (OHTAC) OHTAC Recommendation: Clinical Utility of Vitamin D Testing Revised June 2010.
- Pittas, A.G. MD; Chung, M. MPH et. al Systematic Review: Vitamin D and Cardiometabolic Outcomes. *Ann Intern Med* March 2, 2010 152:5 307-314.
- Ross, A. Catharine, Christine L. Taylor, Ann L. Yaktin, and Heather B. Del Valle, Editors [Dietary Reference Intakes for Calcium and Vitamin D](#); Institute of Medicine, National Academies of Sciences, Washington, DC Nov. 2010. (Key Reference for Community-wide Guideline for Vitamin D Recommendations)
- Stechschulte, S. BA, Kirsner, R MD PhD et al. Vitamin D: Bone and Beyond, Rationale and Recommendations for Supplementation. *The American Journal of Medicine* Sept. 2009 122:9 793-802.
- Wagner C MD, Greer, F MD and the Section on Breastfeeding and Committee on Nutrition Prevention of Rickets and Vitamin D Deficiency in Infants, Children and Adolescents. *American Academy of Pediatrics Clinical Report Pediatrics* November 2008 122:5 1142-1152.
- Williams S MD, Malatesta K BS, Norris K MD Vitamin D and Chronic Kidney Disease. *Ethn Dis* 2009 19(4 Suppl5): S5-8, 11.
- World Health Organization International Agency for Research on Cancer (IARC) Vitamin D and Cancer/A Report of the IARC World Health Organization Press, Geneva Switzerland 2008.